

ADMISSION SHEET



Registration Details :

Admission No : IP26-00006647 Admit Date : 25-Jun-2026 Admit Time : 08:14 PM UHID : LBH-00112796

Patient Details :

Patient Name : Master ARJUN KARTHIKEYA ATMAKURI Age : 1 Y 4 M 12 D
Guardian : Mr SRINIVAS ATMAKURI DOB : 13-02-2025 12:49 PM
Gender : Male Religion :
Occupation : Martial Status :
Address (H) : plot no:128 Malakpet Hyderabad Telangana Phone No : 8008307470/ 7702537184
INDIA 500036 E-mail : sreenivas/atmakuri@gmail.com

Admission Details :

Bed Type : DAY CARE Bed No : ER01 Ward Name : GF -EMERGENCY
Room No : ER01 Admission Type : First Visit

Contact Details :

Name : Mr SRINIVAS ATMAKURI Relationship : Father
Contact Address : plot no:128 Malakpet Hyderabad Telangana Phone No : 8008307470
INDIA 500036


Signature


Doctor Details :

Doctor Name : Dr. SINDHURA MUNUKUNTLA Specialisation : GENERAL PEDIATRICS
Referral Doctor : Self. Phone No :
Co-Consultant :


Payment Details :

Payment Mode : DC/CC Card Deposit Amount : 10000.00
Payor Name : MEDI ASSIST INSURANCE TPA PVT LTD

ACTIVITY RECORD FOR BILLING

LBH-00112796 IP26-00006647
 Name: Master ARJUN KARTHIKEYA
 13-02-2025 1 Y 4 M 12 D (M)
 Dr. SINDHURA MUNUKUNTLA
 UHID No  Consultant : _____ Dept : _____
 Date of Admission : _____ Date of Discharge : _____ Time: _____
 Room / Bed No : _____ Ward : _____ Suggested Billable bed type : _____

WARD TRANSFERS

Date	Time	From	To	Signature of Nurse
25/6/26	8:40 pm	ER	212	

Cross Consultation Visit

	Doctors Name	Date	Order No.	Signature
1.				
2.				
3.				
4.				
5.				
6.				
7.				
8.				
9.				
10.				

Ref.No. F/IN/PR/10



**PEDIATRIC IN-PATIENT
MEDICAL RECORD**

LBH-00112796 IP26-00006647
Master ARJUN KARTHIKEYA
13-02-2025 1 Y 4 M 12 D (M)
Dr. SINDHURA MUNUKUNTLA



Patient Name : Arjun Karthikeya

Patient ID# : _____

Consultant : Dr. Sindhura M

Final Diagnosis : Acute Febrile illness (D3) [Pharyngitis]
 Dehydration Adenovirus

Pediatric

LBH-00112796
Master ARJUN KARTHIKEYA
13-02-2025 1 Y 4 M 12 D (M)
Dr. SINDHURA MUNUKUNTLA

Physical Examination

Name: _____

Informant: Parents

Age/Sex: _____

Reliability: Good.

Chief Presenting Complaints & Duration (Chronologically):

Chf Fever 7-2 days.
cold 7-2 days.
Vomiting since 2 days
Decreased oral intake since 1 day

History of present illness :

Chf Fever - high grade (max - 103F)
- intermittent, subsided on taking medication
- but recurred,
- not a/w rash
- No similar complaints in the family
Chf cold - a/w running nose, nose block.
Sneezing.
Not a/w fast breathing / ~~stridor~~

Oral intake ↓

last urine - 3hrs ago.

Vomiting started 2 days back
multiple episodes, non-bilious, non-projectile
contains food & water.

Baby has decreased oral intake since 1 day

Pediatric Multiorgan History & Physical Examination

LBH-00112796 IP26-00006647
Master ARJUN KARTHIKEYA
13-02-2025 1 Y 4 M 12 D (M)
Dr. SINDHURA MUNUKUNTALA




Past History : (Including details of any previous investigation or treatment)

Blank lined area for Past History.

Birth & Neonatal History :

FTWSCS (Head not engaged)
Bwt - 2850, CIAB, NONICU
admission, NONNJ



Birth & Socio Economic History :

About Father : _____

About Mother : _____

Any additional Information : _____

Developmental History :

- Walks can support
- Can speak 10-12 words.
- 0

Immunization History :

immunised upto 1 year, upto date.

Pediatric Multiorgan History & Physical Examination

Anthropometry

Head Circum (cms) _____ (Centile _____) Height (cm) : _____ (Centile _____)
 Weight (kgs) 14.2 (Centile _____)

On Examination :

Temperature : 102 F Pulse Rate: 180/min Description _____
 B.P. CFT < 3sec SPO2 98% at _____
 Resp. rate and type of breathing : 30/min

Rash NO Signs of rehydration
 Lymphadenopathy NO
 Oedema : NO

Respiratory system :

Inspection (any s/o distress) : _____
 Air entry & breath sounds : NVBS+, flat, No added son
 Any addes sounds : _____
 Relevant data from outside (Chest X-Ray, ABG, etc.,) —

Cardiovasclular System :

Inspection of procordium : _____
 Heart Sounds : S1S2+, NO murmurs
 Any murmur : _____
 Relevant date from outside (Chest X-Ray, ECG, ECHO, Etc.,) _____

Per Abdomen :

Inspection ⊕ (N) shape
 Palpation : soft
 Auscultation : BS+
 Spine: N External Genitelia : h
 Relevant data from outside (CT, USG etc.,) _____

Pediatric Multiorgan History & Physical Examination

Central Nervous System :

Level of Consciousness : AVPU/GCS Score :

15/15

Cranial Nerves :

Motor System :

Nutrition :

Tone :

Power

Co-ordinator :

Posture :

Involuntary Movements :

Reflexes :

DTR

Superficials :

Plantars

Sensory System :

Bladder / Bowel :

Clinical Summary & Diagnostic :

Acute febrile illness + Dehydration

Pediatric Multiorgan History & Physical Examination

LBM-00112796 IP26-00006647
Master ARJUN KARTHIKEYA
13-02-2025 1 Y 4 M 12 D (M)
Dr. SINDHURA MUNUKUNTLA


Preventive aspects of the treatment :

Avoid Dehydration

Desired goals of the treatment :

Adequate hydration

Planned Labs :

CBC, CRP, [Blood c/s - collect & keep]
CUE, ~~RSS~~ VBG
Respiratory panel (5 virus)
XRAY Nasopharynx
Extra plain:

Planned Management :

GROGIN PARACETAMOL
IBUPROFEN
2/3rd maintenance IV fluids
- Monitor vitals

Please fill up the following details

1. Name of the Referring Doctor : _____
2. Name of the Referring Hospital : _____
(Including the name of City)
3. Contact number of the Referring Doctor : _____
(Preferring Mobile #)
4. Name of the doctor in Rainbow Team _____ on
whose name the patient is being referred

Doctor's Signature Name _____ Date _____ Time _____

PROGRESS NOTES AND DOCTOR'S ORDER

Date & Time	Progress Notes	Doctor's Order
25/6/26 8:30 PM	SIB Dr. Sindhura DAFI = dehydration signs of dehydration poor oral intake vit - S, Si M-BL-ACC	Plan 10 fluids - separate breast - CE CROCIN
	PIA-Jac Conscious Anxat - Petrified rally w/rt O/C appearance dry - muc Bluept	9 DUKOFEN 500 Monitor vitals of medicine DR. SINDHURA
26/6/26 12:25 AM	Case disc Dr. Sindhura WBC - 12,200 Neutrophils - 62%	send Blood CC Start CEFTRIAXONE AB-Monitahi AM
	B-100	

PROGRESS NOTES AND DOCTOR'S ORDER

Date & Time	Progress Notes	Doctor's Order
26/6/26	c/s/hy Dr. Sindhu M	
9:00 AM	AFI C dehydration Covid (+ve)	
	fever spikes ⊕	<u>Plan</u>
	oral intake - poor.	✓ Cont ceftriaxone
	vitals - stable.	✓ Metatop Nasal spray 4 puff HS.
	R/S - BIL AE ⊕ PIA - soft, NT	✓ Trace B/Cfs Adeno viral.
		✓ Monitor vitals Noted by Riva 26/6/26
		P. Sundar Dr. Sindhu M.

LBH-00112796 IP26-00006647
 Master ARJUN KARTHIKEYA
 13-02-2025 1 Y 4 M 12 D (M)
 Dr. SINDHURA MUNUKUNTLA



MEDICATION RECONCILIATION FORM

Drug Allergies: Not known any Drug Allergies

**Medication Reconciliation will be done at the time of admission and also whenever there is change in the treating team or shifting from one unit to another unit.
 (Example: at the time of admission shifting from ICU to Ward, or Ward to ICUs)**

Shifting From: ER Shifted to: ward

S.No	MEDICATION NAME (GENERIC NAME CAPITAL LETTERS)	DOSE (mg, mcg)	ROUTE (PO, NG, SC, IV)	FREQUENCY	LAST DOSE Date / Time	ON ADMISSION / SHIFTING
1						<input type="checkbox"/> C <input type="checkbox"/> DC
2						<input type="checkbox"/> C <input type="checkbox"/> DC
3						<input type="checkbox"/> C <input type="checkbox"/> DC
4						<input type="checkbox"/> C <input type="checkbox"/> DC
5						<input type="checkbox"/> C <input type="checkbox"/> DC
6						<input type="checkbox"/> C <input type="checkbox"/> DC
7						<input type="checkbox"/> C <input type="checkbox"/> DC
8						<input type="checkbox"/> C <input type="checkbox"/> DC
9						<input type="checkbox"/> C <input type="checkbox"/> DC
10						<input type="checkbox"/> C <input type="checkbox"/> DC

* C- Continue, DC - Discontinue

MEDICATION HISTORY RECORDED / VERIFIED BY


Doctor Name & Signature : Dr. Prasanti

Date & Time : 25/6/26 @ 7:30 Pm

Nurse Name & Signature: Anupam

Date & Time : 25/6/26 @ 7:20 Pm

Docu. No. : RCH / FRM / GENERAL / 090





14 kg

DRUG CHART

Date of Admission: Drug Allergies: Not known any Drug Allergies

FOR THE SAFETY OF THE PATIENT

- GENERAL** - Ensure that all patient details are entered above. ONLY A DOCTOR SHALL WRITE MEDICATION ORDERS.
- DOCTOR** - Please use only approved abbreviations (refer to Hospital's approved list of abbreviations).
 - Use approved pharmaceutical names, BLOCK LETTERS, metric dosage. English instructions.
 - Any changes in drug therapy must be ordered by a NEW PRESCRIPTION. Do not alter existing instructions.
 - Discontinue a drug by drawing a line **I** through it and a similar line through subsequent recording panels.
 - The date and time of stopping the drug along with the doctors name and sign must be mentioned.
 - Only one chart should be in use at any one time. When the chart is full, a new supplement can be kept within this drug sheet folder.
- NURSES** - Nurses must follow strictly the FIVE RIGHTS before administration of medication.
 1) Right Patient 2) Right Drug 3) Right Dosage 4) Right Route 5) Right Time
 - AVOID TAKING VERBAL ORDERS. NO VERBAL ORDERS FOR HIGH RISK/HIGH ALERT MEDICINES (EXCEPT FIRST DOSE OF EPINEPHRINE DURING CPR). Follow Hospitals's Verbal Order Policy.

SOS / PRN (As Required Medication)

DRUG : SYP. PARACETAMOL				Date Time															
Dose	Route	Frequency	Start Date																
240 4.5ml (PO)		SOS	25/6																
Doctor's Signature		Valid Period	Pharm.																
Ruth																			
Additional Instructions:																			
SYP. CROCIN DS (240/5)																			

DRUG : SYP. IBUGESIC				Date Time															
Dose	Route	Frequency	Start Date																
3.5ml	PO	SOS	25/6																
Doctor's Signature		Valid Period	Pharm.																
Ruth																			
Additional Instructions:																			
C 100/5 SYP. IBUPROFEN																			

DRUG :				Date Time															
Dose	Route	Frequency	Start Date																
Doctor's Signature		Valid Period	Pharm.																
Additional Instructions:																			

VERIFIED BY: NAME Signature

REGULAR PRESCRIPTIONS

Weight. 14.2kg Ward.



DRUG : Sy P. XYRAL				Date Time	25/6																
Dose	Route	Frequency	Start Date																		
2.5ml	oral	bedtime	25/6																		
Name & Signature of the Doctor Starting the Drugs: B. Sreelakshmi				10PM 25/6																	
Additional Instructions: LEVOCETRIZINE (2.5mg/5ml)																					
Daily Doctor's Endorsement by a Sign																					
DRUG : 90% CEFTRIAZONE				Date Time	25/6																
Dose	Route	Frequency	Start Date																		
1.4gm	IV	OD	26/6																		
Name & Signature of the Doctor Starting the Drugs: B. Sreelakshmi				10PM 11AM 25/6																	
Additional Instructions: → dilute in 30ml NS and give over 2 hours																					
Daily Doctor's Endorsement by a Sign																					
DRUG : Metatop Nasal Spray				Date Time																	
Dose	Route	Frequency	Start Date																		
1 Puff	Nasal	HS.	26/6																		
Name & Signature of the Doctor Starting the Drugs: A. Sreelakshmi																					
Additional Instructions:																					
Daily Doctor's Endorsement by a Sign																					
DRUG :				Date Time																	
Dose	Route	Frequency	Start Date																		
Name & Signature of the Doctor Starting the Drugs:																					
Additional Instructions:																					
Daily Doctor's Endorsement by a Sign																					

wt - 14.15kg

EMERGENCY ROOM TRIAGE FORM

Patient's Name : Arjun Karthikeya Age : 1.5 year Gender: Male Female

Date : 25/6/20 Time of Arrival : 7:04 PM

Allergies: No Yes Food Medications Blood Transfusion Other (Specify): Not known

Source of Information: Parents Others (Specify)

Mode of Arrival: Ambulatory Wheelchair Ambulance

Initial Vital Signs: Temp: 38.5 PR: 203b/m BP: RR: SpO₂: 100%

Chief Complaints: C/O Fever since 2 days

INITIAL PHYSIOLOGICAL CATEGORIZATION Appearance <input checked="" type="checkbox"/> Normal <input type="checkbox"/> Sick Looking Work of Breathing <input checked="" type="checkbox"/> Normal <input type="checkbox"/> Increased <input type="checkbox"/> Decreased <input type="checkbox"/> Gasping / Apnea Circulation / Colour <input checked="" type="checkbox"/> Normal <input type="checkbox"/> Abnormal <input type="checkbox"/> Bleeding		INITIAL PHYSIOLOGICAL STATUS <input checked="" type="checkbox"/> Stable <input type="checkbox"/> Unstable : <input type="checkbox"/> Not - Life - Threatening <input type="checkbox"/> Life -Threatening
---	--	---

Triage Classification	CTAS
<input type="checkbox"/> Level 1 : Resuscitation	<input type="checkbox"/> Immediate
<input type="checkbox"/> Level 2 : EMERGENT : Life or limb threatening	<input type="checkbox"/> < 15 min
<input type="checkbox"/> Level 3 : URGENT : Significant illness / injury with potential to become life or limb threatening	<input type="checkbox"/> 30 min
<input type="checkbox"/> Level 4 : LESS URGENT : Significant illness but not life threatening	<input checked="" type="checkbox"/> 60 min
<input type="checkbox"/> Level 5 : NON - URGENT : May receive care when convenient	<input type="checkbox"/> 120 min

NOTE : All immunocompromised children and preterm babies to be considered Level 2.
 All Children less than 2 years age with high fever to be considered Level 3.
 * CTAS - Canadian Triage and Acuity Scale

Signature of Parent / Guardian
 Triage Completion Time : 7:06 PM

Communicable Disease Triage Screening

PART A. The following questions should be asked to all patients at the initial screening:

- Have you had fever (elevated temperature) in the past 2 weeks Yes No
- Have you had cough or a rash in the past 2 weeks Yes No
- Have you had shortness of breath or difficulty breathing in the past 2 weeks Yes No

PART B. For patients reporting fever and respiratory/rash symptoms: Not applicable

- Have you travelled outside the INDIA? or had close contact with someone who has recently travelled outside the INDIA, in the past two weeks? Yes No
 If yes, State Location:
- Are your parents / close contacts at home is/a healthcare worker? {please encircle the choices} (e.g., nurse, physician, ancillary services personnel, allied health services personnel, hospital volunteer, or laboratory worker, others) who has had a recent exposure to an individual with a highly communicable disease or unexplained, severe febrile respiratory or rash disease? Yes No

PART C. A positive communicable disease triage screening is considered for any patient who meets one of the two following criteria:

- Any patient with Fever / Rash / Vesicles / Discharge from Eyes and Cough
- Any patient with fever and respiratory symptoms who answered "YES" to any of the questions on epidemiologic risk factors in "PART B" of the triage screening above.

PART D. ACTION / INTERVENTION: (for positive suspected communicable disease triage screening)

- Patients should be immediately isolated in a negative pressure room or a single room (as appropriate) for pending evaluation.
- The patient should be given a surgical mask immediately, if not already wearing one.
- Both patient and triage staff should perform hand hygiene.
- The staff should use PPE (as appropriate).

Name of Triage Nurse : Prabin
 Date & Time : 25/6/20 @ 7:06 PM
 Docu. No. : RCH /FRM / CLINICAL / 085

Signature of Triage Nurse : [Signature]

NURSING INITIAL ASSESSMENT IN EMERGENCY ROOM

Date: 25/6/26 Time of arrival: 7:04 PM

Chief Complaints: @ @ Fever since 2 days RBS:

Height: Weight: BMI: Head Circumference (<2 years)

Allergies: Yes No Medications Blood Transfusion Food Other:

If yes, identify

Pain Screening: Yes No If Yes, Pain Score: 10 Pain Tool Used: N Pass FLACC Wong Baker

Character Location Frequency Duration

RISK FOR FALL:

- If patient is < 6 years
tick below fall risk intervention directly
- If Patient is > 6 years
Assess the below parameters

History of Falling: within past 3 months Yes No

Ambulatory Aids:

- Wheelchair Yes No
- Uses furniture for support Yes No

Gait/Transferring:

- Bedrest / immobile Yes No
- Weak Yes No
- Impaired Yes No

Mental Status: Forgets limitations Yes No

IF YES FOR ANY CATEGORY = RISK FOR FALLING

Fall Risk Intervention:

- Escort while ambulating
- Assist Patient
- Educate patient and family on fall precautions/prevention

Functional Screening: No Abnormalities Detected

- Mobility Problem
- Walking Problem
- Developmental Delay
- Musculoskeletal Congenital Abnormality

Inform consultant for positive criteria

.....

.....

Nutritional Screening: No Abnormalities Detected

- Underweight
- Overweight
- Feeding Problem
- Special diet
- Special feeding method

Inform consultant for positive criteria

Psychological Screening: No Significant Findings

Unusual concerns about patient's Psychological Status: Yes No

If Yes Consultant Notified: (Date/Time):

Social History: Lives With Family

Siblings in household Yes No (if yes How Many?)

Time of Initial assessment completed by ER Nurse: 7:06 PM

Nursing Notes (Including Labs / Medications / Other Care):

Time	Nursing Notes
	→ Assessed the pt condition
	→ checked the pt vitals
	→

Samples collected by:

Time:

Samples sent by:

Apurba

Time:

8:20pm

Medication given in ER:

Date / Time	Medication	Route	Dosage & Instructions	Doctor Sign	Nurse Sign 1
6:30pm	Crocin	PO	6 ml		<i>[Signature]</i>

Condition of patient at time of shift - out :	Details of Shift - out
HR: <u>141</u> BP: CFT:	Shift - out from ER to: <u>212</u>
RR: <u>29</u> SPO ₂ : <u>100%</u>	Time of Shift - out: <u>8:40pm</u>
GCS: Temperature: <u>99</u>	Handover given to: <u>Moufushi</u>
Pain Score: <u>0</u>	(Nurse's Name)
Repeat RBS (if applicable):	

Tick as applicable: MLC LAMA BROUGHT DEAD


Procedures done with details (if any):

Name of the Nurse: Brabi'r

Signature of the Nurse: *[Signature]*

Date & Time: 25/6/26 @ 7:06

PATIENT TRANSFER FORM

Patient Name & UHID No. LBH-00112796 IP26-00006647 Master ARJUN KARTHIKEYA 13-02-2025 1 Y 4 M 12 D (M) Dr. SINDHURA MUNUKUNTLA 		Date & Time of Admission 25/6/20	Date & Time of Transfer Order 25/6/20
		Transfer Ordered by Dr. Prasanti	Reason for Transfer Admission
From Unit ER	To Unit 212	Information to Attendant Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>	
Number of Sheets in Clinical File 21	Number of Imaging Films 1	Personal belongings including clinical documents. If any handed over to attendant Yes <input type="checkbox"/> No <input type="checkbox"/> If yes, what ?	
Medications / Consumables / Surgicals / Hand over			
Sl.No.	Item Name	Quantity	
1.			
2.			
3.			
4.			
5.			
Shifting Summary / Notes Written by Doctor : Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>			
Name & Signature of Person who is Transferring Anupam		Name of Person Ordered Transfer Dr. Prasanti	
Patient & Clinical Records Received by : Moulishi			
Date & Time of Patient Received : 25/6/20 @ 10:pm.			

If the transfer order time & Completion time is more than 30 minutes, please tick the reason mentioned below :

- Unavailable Bed Nurse not Available Available Bed not ready

LBH-00112796 IP26-00006647

Master ARJUN KARTHIKEYA

13-02-2025 1 Y 4 M 13 D (M)

Dr. SINDHURA MUNUKUNTLA



212



RESULT SHEET

Date	25/6/26				
Time					
Hb	9.4				
PCV	28.8				
RBC	5.04				
WBC	10.20				
N/L	61.9/31.5				
Platelets	340				
CRP	5.0				
ESR					
PCT					
RBS					
Na					
K					
Cl					
Ca/Mg					
Phosphate					
Urea					
Creatinine					
ALP					
SGPT					
SGOT					
T.Bill/Conj					
T.Protein					
S.Albumin					
S.Globulin					
A/G Ratio					
Uric Acid					
S.Amylase					
Sr.Lipase					
Blood Lactate					
S.Cholesterol					
PT/INR					
APTT					
CSF Protein / Sugar					
Cells					
N/L					

Date	26/6/26					
Time						
CUE - Alb	.					
CUE - Sugar						
CUE - Ketones	present					
CUE - PUS Cells	4-6					
CUE - RBC Cells	Nil					
CUE						
Epithelial cells -	1-2					
Leucocyte -	Negative					
Stool Pus Cell						
OVA / Cyst						
Occult Blood						

Culture and Sensitivities :

.....

.....

.....

Radiology : USG :

 X-Ray :

 ECHO :

 CT :

 MRI :

 Others (ECG, Contrast Studies etc.,) :

EARLY WARNING SCORE: CHILDREN'S UNIT

Date : 25/6..... Time: 10pm	2	3	4	5	6	7	8	9	10	11	12
Doctor / Nurse / Family Concern?	AM	AM	AM	AM	AM	AM	AM	AM	AM	AM	AM
Temperature (F)	98.1 F	99.5 F	101.8 F	100.9 F	99.9	98.4 F	98.1 F				
Heart Rate (bpm) and Blood Pressure (mmHg) *	145b/m	135b/m					150b/m				
Resp. Rate (bpm) (Over 1 Minute) *	35b/m	35b/m					38b/m				
Receiving O ₂ (l/min) / O ₂ Saturations (%)	2a/liter / 96%	0.1 liter / 98%					0.1 liter / 99%				
Conscious Level	Normal	Normal					Normal				
GCS *											
TOTAL SCORE											
Number of shaded boxes	0	0					0				
Pain Score	0	0					0				
Observer's Initials	AS	AS					AS				

ACTIONS

Score 1 : Continue normal observation by staff nurse
 Score 2 : Shift in charge nurse to be informed and continue hourly observations
 Score 3 : Shift in charge AND ER doctor/Floor Registrar to see and half hourly to hourly Observation to continue.
 Score 4 : Shift in charge AND treating consultant(till 8 PM) or On call night duty consultant to see
 Score 5 & 6 : Shift in charge AND PICU fellow or PICU consultant to be informed.

NB: Scores 3 should be recorded overleaf

* NB: If GCS is below 12 or the Oxygen requirement is >3 Lit./min. , then irrespective of rest of the score, the Nurse MUST inform the PICU team.

CHILDREN'S OBSERVATION and EARLY WARNING SCORING TOOL

INSTRUCTIONS:

- The paediatric Early Warning Score i) seeks to identify the abnormal physiological finding seen during serious childhood illnesses and ii) offers a method to interpret such physiological derangements with clearly defined actions, ensuring that suitably experienced staff are involved with the care of the sickest children.
- The Early Warning Score does not replace clinical experience and acumen and should not be relied upon for such purpose.
- 6 clinical parameters are assessed and recorded as part of the child's routine clinical observation, providing a Early Warning Score between 0-6 (Higher Early Warning Score are seen in sicker children)
- Detailed actions are described according to increasing Early Warning Score.
- Some children with complex medical needs e.g. cyanotic heart disease may require modification to their trigger thresholds/ action plan- this should follow discussion with senior colleagues.
- Any Early Warning Score of 3 or above should be recorded below with details of any subsequent action initiated

Record Details when EARLY WARNING SCORE > 3			Record Time of Review and Plan		
Date	Time	Early Warning Score	Date	Time	Name

- If at any time additional help is required, call help – regardless of the Early Warning Score!
- Following a Early Warning Score assessment, senior help may be required

The SBAR communication tool (situation, background, assessment, recommendations) is a helpful mnemonic that can be used to describe a child's clinical condition to a colleague.

I	IDENTITY: I am (name), a nurse on ward (X). I am calling about (child X)
S	SITUATION: I am calling because I am concerned that ... (e.g. BP is low/high, pulse is XXX, Temperature is XX, Early Warning Score is XX)
B	BACK GROUND: Child (X) was admitted on (XX date) with (e.g. respiratory infection). They have had (X operation/ procedure/ investigation). Child (X)'s condition has changed in the last (XX mins). Their last set of observations were (XXX). The child's normal condition is ... (e.g. alert/ drowsy/ confused, pain free)
A	ASSESSMENT: I think the problem is (XXX) and I have ... (e.g. given O2/ analgesia, stopped the infusion), OR I am not sure what the problem is but child (X) is deteriorating, OR I don't know what's wrong but I am really worried.
R	RECOMMENDATION: I need you to ... come to see the child in the next (XX mins) AND is there anything I need to do in the meantime? (e.g. stop the fluid/ repeat observation)

CHILDREN'S OBSERVATION and EARLY WARNING SCORING TOOL

INSTRUCTIONS:

- The paediatric Early Warning Score i) seeks to identify the abnormal physiological finding seen during serious childhood illnesses and ii) offers a method to interpret such physiological derangements with clearly defined actions, ensuring that suitably experienced staff are involved with the care of the sickest children.
- The Early Warning Score does not replace clinical experience and acumen and should not be relied upon for such purpose.
- 6 clinical parameters are assessed and recorded as part of the child's routine clinical observation, providing a Early Warning Score between 0-6 (Higher Early Warning Score are seen in sicker children)
- Detailed actions are described according to increasing Early Warning Score.
- Some children with complex medical needs e.g. cyanotic heart disease may require modification to their trigger thresholds/ action plan- this should follow discussion with senior colleagues.
- Any Early Warning Score of 3 or above should be recorded below with details of any subsequent action initiated

Record Details when EARLY WARNING SCORE > 3			Record Time of Review and Plan		
Date	Time	Early Warning Score	Date	Time	Name

- If at any time additional help is required, call help – regardless of the Early Warning Score!
- Following a Early Warning Score assessment, senior help may be required

The SBAR communication tool (situation, background, assessment, recommendations) is a helpful mnemonic that can be used to describe a child's clinical condition to a colleague.

I	IDENTITY: I am (name), a nurse on ward (X). I am calling about (child X)
S	SITUATION : I am calling because I am concerned that ... (e.g. BP is low/high, pulse is XXX, Temperature is XX, Early Warning Score is XX)
B	BACK GROUND : Child (X) was admitted on (XX date) with (e.g. respiratory infection). They have had (X operation/ procedure/ investigation). Child (X)'s condition has changed in the last (XX mins). Their last set of observations were (XXX). The child's normal condition is ... (e.g. alert/ drowsy/ confused, pain free)
A	ASSESSMENT : I think the problem is (XXX) and I have ...(e.g. given O2/ analgesia, stopped the infusion), OR I am not sure what the problem is but child (X) is deteriorating, OR I don't know what's wrong but I am really worried.
R	RECOMMENDATION : I need you to ... come to see the child in the next (XX mins) AND I s there anything I need to do in the meantime ? (e.g. stop the fluid/ repeat observation)

CHILDREN'S OBSERVATION and EARLY WARNING SCORING TOOL

INSTRUCTIONS:

- The paediatric Early Warning Score i) seeks to identify the abnormal physiological finding seen during serious childhood illnesses and ii) offers a method to interpret such physiological derangements with clearly defined actions, ensuring that suitably experienced staff are involved with the care of the sickest children.
- The Early Warning Score does not replace clinical experience and acumen and should not be relied upon for such purpose.
- 6 clinical parameters are assessed and recorded as part of the child's routine clinical observation, providing a Early Warning Score between 0-6 (Higher Early Warning Score are seen in sicker children)
- Detailed actions are described according to increasing Early Warning Score.
- Some children with complex medical needs e.g. cyanotic heart disease may require modification to their trigger thresholds/ action plan- this should follow discussion with senior colleagues.
- Any Early Warning Score of 3 or above should be recorded below with details of any subsequent action initiated

Record Details when EARLY WARNING SCORE > 3			Record Time of Review and Plan		
Date	Time	Early Warning Score	Date	Time	Name

- If at any time additional help is required, call help – regardless of the Early Warning Score!
- Following a Early Warning Score assessment, senior help may be required

The SBAR communication tool (situation, background, assessment, recommendations) is a helpful mnemonic that can be used to describe a child's clinical condition to a colleague.

I	IDENTITY: I am (name), a nurse on ward (X). I am calling about (child X)
S	SITUATION : I am calling because I am concerned that ... (e.g. BP is low/high, pulse is XXX, Temperature is XX, Early Warning Score is XX)
B	BACK GROUND : Child (X) was admitted on (XX date) with (e.g. respiratory infection). They have had (X operation/ procedure/ investigation). Child (X)'s condition has changed in the last (XX mins). Their last set of observations were (XXX). The child's normal condition is ... (e.g. alert/ drowsy/ confused, pain free)
A	ASSESSMENT : I think the problem is (XXX) and I have ... (e.g. given O2/ analgesia, stopped the infusion), OR I am not sure what the problem is but child (X) is deteriorating, OR I don't know what's wrong but I am really worried.
R	RECOMMENDATION : I need you to ... come to see the child in the next (XX mins) AND I s there anything I need to do in the meantime ? (e.g. stop the fluid/ repeat observation)

LBM-00112796 IP26-00006647
 Master ARJUN KARTHIKEYA
 13-02-2025 1 Y 4 M 13 D (M)
 Dr. SINDHURA MUNUKUNTLA



FLUID CHART

Sheet No. :

1. All measurements in ml.
2. Add up each column separately. Make additions across the page to obtain 24 hrs. total of intake and output.

Date	Time	Nature of Fluid	Intake			Output					IV Site Thrombophlebitis Score	Sign. Nurse	
			Mouth	I.V	N.G	NG	Diarrhoea	Vomit	Drainage	Urine			
	08:00 am												
	09:00 am												
	10:00 am												
	11:00 am												
	12:00 pm												
	01:00 pm												
Total Intake :						Total Output :							
	02:00 pm												
	03:00 pm												
	04:00 pm												
	05:00 pm												
	06:00 pm												
	07:00 pm												
Total Intake :						Total Output :							
25/6/26	08:00 pm												
	09:00 pm												
	10:00 pm				33ml								
	11:00 pm				33ml								
	12:00 am				33ml								
	01:00 am				33ml								
Total Intake :						Total Output :							
26/6/26	02:00 am				33ml								
	03:00 am				33ml								
	04:00 am				33ml								
	05:00 am				33ml								
	06:00 am				33ml								
	07:00 am				33ml								
Total Intake :						Total Output :							

Total 24 hrs. Intake

Total 24 hrs. Output

LBH-00112798 1P26-00006647
 Master ARJUN KARTHIKEYA
 13-02-2026 1 Y 4 M 13 D (M)
 Dr. SINDHURA MUNUKUNTLA



FLUID CHART

Sheet No. :

1. All measurements in ml.
2. Add up each column separately. Make additions across the page to obtain 24 hrs. total of intake and output.

Date	Time	Nature of Fluid	Intake			Output					IV Site Thrombo-phlebitis Score	Sign. Nurse		
			Mouth	I.V	N.G	NG	Diarrhoea	Vomit	Drainage	Urine				
	08:00 am													
	09:00 am													
	10:00 am													
	11:00 am													
	12:00 pm													
	01:00 pm													
Total Intake :						Total Output :								
	02:00 pm													
	03:00 pm													
	04:00 pm													
	05:00 pm													
	06:00 pm													
	07:00 pm													
Total Intake :						Total Output :								
	08:00 pm													
	09:00 pm													
	10:00 pm													
	11:00 pm													
	12:00 am													
	01:00 am													
Total Intake :						Total Output :								
	02:00 am													
	03:00 am													
	04:00 am													
	05:00 am													
	06:00 am													
	07:00 am													
Total Intake :						Total Output :								

Total 24 hrs. Intake	
-----------------------------	--

Total 24 hrs. Output	
-----------------------------	--

Patient Sticker



FLUID CHART

Sheet No. :

1. All measurements in ml.
2. Add up each column separately. Make additions across the page to obtain 24 hrs. total of intake and output.

Intake						Output					IV Site Thrombo- phlebitis Score	Sign. Nurse	
Date	Time	Nature of Fluid	Route			NG	Diarrhoea	Vomit	Drainage	Urine			
			Mouth	I.V	N.G								
	08:00 am												
	09:00 am												
	10:00 am												
	11:00 am												
	12:00 pm												
	01:00 pm												
Total Intake :						Total Output :							
	02:00 pm												
	03:00 pm												
	04:00 pm												
	05:00 pm												
	06:00 pm												
	07:00 pm												
Total Intake :						Total Output :							
	08:00 pm												
	09:00 pm												
	10:00 pm												
	11:00 pm												
	12:00 am												
	01:00 am												
Total Intake :						Total Output :							
	02:00 am												
	03:00 am												
	04:00 am												
	05:00 am												
	06:00 am												
	07:00 am												
Total Intake :						Total Output :							

Total 24 hrs. Intake	
-----------------------------	--

Total 24 hrs. Output	
-----------------------------	--

Patient Sticker



FLUID CHART

Sheet No. :

1. All measurements in ml.
2. Add up each column separately. Make additions across the page to obtain 24 hrs. total of intake and output.

		Intake				Output						
Date	Time	Nature of Fluid	Route			NG	Diarrhoea	Vomit	Drainage	Urine	IV Site Thrombo-phlebitis Score	Sign. Nurse
			Mouth	I.V	N.G							
	08:00 am											
	09:00 am											
	10:00 am											
	11:00 am											
	12:00 pm											
	01:00 pm											
	Total Intake :						Total Output :					
	02:00 pm											
	03:00 pm											
	04:00 pm											
	05:00 pm											
	06:00 pm											
	07:00 pm											
Total Intake :						Total Output :						
	08:00 pm											
	09:00 pm											
	10:00 pm											
	11:00 pm											
	12:00 am											
	01:00 am											
Total Intake :						Total Output :						
	02:00 am											
	03:00 am											
	04:00 am											
	05:00 am											
	06:00 am											
	07:00 am											
Total Intake :						Total Output :						

Total 24 hrs. Intake	
-----------------------------	--

Total 24 hrs. Output	
-----------------------------	--

LBH-00112796 IP26-00006647
 Master ARJUN KARTHIKEYA
 13-02-2025 1 Y 4 M 13 D (M)
 Dr. SINDHURA MUNUKUNTLA



NURSING CARE RECORD



Date: 25/6/26

- Goals**
- Maintain Airway and Oxygenation
 - Relieve Pain & Discomfort
 - Maintain Fluid Balance
 - Improve Activity Tolerance
 - Maintain Good Nutritional Status
 - Maintain Skin Integrity
 - Maintain Personal Hygiene
 - Prevent Infection
 - Meet Elimination Needs
 - Ensure Safety
 - Early Ambulation Reduce Anxiety
 - Patient & Family Education
 - Identify Potential Complications
 - Any Others. Specify.....

	Time	Plan of Care	Time	Implementation	Evaluation	Re-Assessment	Nurse Name & Signature
Morning							
Afternoon							
Night		<p>Assess the Baby general condition</p> <p>- maintain Ito chart</p> <p>- Administration medication as per doctor advice</p>		<p>assessed Baby general condition</p> <p>checked vital & recorded</p> <p>Administered medication as per doctor advice</p> <p>maintained Ito chart</p>	<p>pt is stable</p>	<p>see checked vital</p>	

Night
Docu

Patient Sticker

NURSING CARE RECORD



Date:

Goals

- Maintain Airway and Oxygenation
- Maintain Personal Hygiene
- Identify Potential Complications
- Relieve Pain & Discomfort
- Prevent Infection
- Any Others. Specify.....
- Maintain Fluid Balance
- Meet Elimination Needs
- Improve Activity Tolerance
- Ensure Safety
- Maintain Good Nutritional Status
- Early Ambulation Reduce Anxiety
- Maintain Skin Integrity
- Patient & Family Education

	Time	Plan of Care	Time	Implementation	Evaluation	Re-Assessment	Nurse Name & Signature
Morning							
Afternoon							
Night							

Morning
Afternoon
Night

BRADEN 'Q' SCALE

Date : 23/6
 Time : 12:1

Mobility	1. Completely immobile: Does not make even slight changes in body or extremity position without assistance.	2. Very limited: Makes occasional slight changes in body or extremity position but unable to completely turn self independently.	3. Slightly limited: Makes frequent through slight changes in body or extremity position independently.	4. No limitations: Makes major and frequent changes in position without assistance.	4		
"Activity The degree of physical activity"	1. Bedfast : Confined to bed	2. Chairfast : Ability to walk severely limited or non-existent. Cannot bear own weight and/or must be assisted into chair or wheelchair."	3. Walks occasionally: Walks occasionally during day, but for very short distances, with or without assistance. Spends majority of each shift in bed or chair.	4. All patients too young to ambulate; OR walks frequently: Walks outside the room at least twice a day and inside room at least once every 2 hours during walking hours.	4		
Sensory Perception	1. Completely limited: Unresponsive (does not moan, flinch or grasp) to painful stimuli due to diminished level of consciousness or sedation. OR, limited ability to feel pain over most of the body surface.	2. Very limited: responds to only painful stimuli, cannot communicate discomfort except by moaning or restlessness; OR, has sensory impairment that limits the ability to feel pain or discomfort over half of body.	3. Slightly limited: Responds to verbal commands, but cannot always communicate discomfort or need to be turned; OR, has some sensory impairment that limits ability to feel pain, or discomfort in one or two extremities.	4. No impairment: Responds to verbal commands. Has no sensory deficit that would limit ability to feel or communicate pain or discomfort.	4		
Moisture Degree to which skin is exposed to moisture	1. Constantly moist: Skin is kept moist almost constantly by perspiration, urine, drainage, etc. Dampness is detected every time patient is moved or turned.	2. Very moist: Skin is often, but not always, moist. Linen must be changed at least every 8 hours.	3. Occasionally moist: Skin is occasionally moist, requiring linen change every 12 hours.	4. Rarely moist: Skin is usually dry, routine diaper changes; linen only requires changing every 24 hours.	4		
FRICITION-SHEAR Friction Occurs when Skin moves against support surfaces Shear Occurs when skin and adjacent bony surface slide across one another	1. Significant problem: Spasticity, contracture, itching, or agitation leads to almost constant thrashing and friction.	2. Problem: Requires moderate to maximum assistance in moving. Complete lifting without sliding against sheets is impossible. Frequently slides down in bed or chair, requiring frequent repositioning with maximum assistance.	3. Potential problem: Moves freely or requires minimum assistance. During a move, skin probably slides to some extent against sheets, chair, restraints, or other devices. Maintains relative good position in chair or bed most of the time but occasionally slides down.	4. No apparent problem: Able to completely lift patient during position change, moves in bed and in chair independently and has sufficient muscle strength to life up completely during move. Maintains good position in bed or chair at all times."	4		
Nutritional Usual food intake pattern	1. Very Poor: NPO/or maintained on clear liquids, or IVs for more than 5 days OR albumin < 2.5 mg/dl OR never eats a complete meal. Rarely eats more than half of any food offered. Protein intake includes only 2 servings or meat or dairy products per day. Takes fluids poorly. Does not take a liquid dietary supplement.	2. Inadequate: Is on liquid diet or tube feedings/TPN, which provides inadequate calories and minerals for age OR albumin < 3 mg/dl OR rarely eats a complete meal and generally eats only about half of any food offered. Protein intake includes only 3 servings of meat or dairy products per day. Occasionally will take a dietary supplement.	3. Adequate: Is on tube feedings or TPN, which provide adequate calories and minerals for age OR eats over half of most meals. Eats a total of 4 servings of protein (meat, dairy products) each day. Occasionally will refuse a meal, but will usually take a supplement if offered.	4. Excellent: Is on a normal diet providing adequate calories for age. For example, eats most of every meal. Never refuses a meal. Usually eats a total of 4 or more servings of mean and dairy products. Occasionally eats between meals. Does not require supplementation.	4		
Tissue Perfusion & Oxygenation	1. Extremely compromised: Hypotensive (MAP < 50 mm Hg; < 40 in a newborn) or the patient does not physiologically tolerate position changes.	2. Compromised: Normotensive oxygen saturation may be < 95%; hemoglobin may be < 10 mg/dl; capillary refill may be > 2 seconds; serum pH is < 7.40.	3. Adequate: Normotensive oxygen saturation may be < 95%; hemoglobin may be < 10 mg/dl; capillary refill may be < 2 seconds; serum pH is normal.	4. Excellent: Normotensive, oxygen saturation > 95%; normal hgb; capillary refill < 2 seconds.	4		

TOTAL SCORE	28		
Evaluator's Name	CL		

Severe Risk : less than 9 | High Risk : 10-12 | Moderate Risk : 13-14 | Mild Risk : 15-18 | Not at Risk: 19-23

Risk Score	Category	Action	Support Surfaces (Please Note: Only required for children who are deemed at risk due to altered mobility, consider occupation therapy referral for advice)
15-18	At Risk	<ul style="list-style-type: none"> • Regular Turning Schedule • Enable as much activity as possible • Protect the heels • Use pressure redistribution surfaces • Manage moisture, friction and shear • Advance to a higher level of risk if other major risk factors are present 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
13-14	Moderate Risk	<ul style="list-style-type: none"> • Use the Same Protocol as for “At Risk” Patients • Position patient at 30 degree lateral incline using foam wedges 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
10-12	High Risk	<ul style="list-style-type: none"> • Follow the same protocol as for “Moderate Risk” Patients • In addition to regular turning schedule • Make small shifts in their position frequently 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
Less than 9	Severe Risk	<ul style="list-style-type: none"> • Use same protocol as for “High Risk” Patients • Add a pressure redistribution surface for patients with severe pain or with additional risk factors. 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay

Patient ID _____

BRADEN 'Q' SCALE



					Date :				
					Time :				
Mobility	1. Completely immobile: Does not make even slight changes in body or extremity position without assistance.	2. Very limited: Makes occasional slight changes in body or extremity position but unable to completely turn self independently.	3. Slightly limited: Makes frequent through slight changes in body or extremity position independently.	4. No limitations: Makes major and frequent changes in position without assistance.					
Activity The degree of physical activity	1. Bedfast : Confined to bed	2. Chairfast : Ability to walk severely limited or non-existent. Cannot bear own weight and/or must be assisted into chair or wheelchair.*	3. Walks occasionally: Walks occasionally during day, but for very short distances, with or without assistance. Spends majority of each shift in bed or chair.	4. All patients too young to ambulate; OR walks frequently: Walks outside the room at least twice a day and inside room at least once every 2 hours during walking hours.					
Sensory Perception	1. Completely limited: Unresponsive (does not moan, flinch or grasp) to painful stimuli due to diminished level of consciousness or sedation, OR, limited ability to feel pain over most of the body surface.	2. Very limited: responds to only painful stimuli, cannot communicate discomfort except by moaning or restlessness; OR, has sensory impairment that limits the ability to feel pain or discomfort over half of body.	3. Slightly limited: Responds to verbal commands, but cannot always communicate discomfort or need to be turned; OR, has some sensory impairment that limits ability to feel pain, or discomfort in one or two extremities.	4. No impairment: Responds to verbal commands. Has no sensory deficit that would limit ability to feel or communicate pain or discomfort.					
Moisture Degree to which skin is exposed to moisture	1. Constantly moist: Skin is kept moist almost constantly by perspiration, urine, drainage, etc. Dampness is detected every time patient is moved or turned.	2. Very moist: Skin is often, but not always, moist. Linen must be changed at least every 8 hours.	3. Occasionally moist: Skin is occasionally moist, requiring linen change every 12 hours.	4. Rarely moist: Skin is usually dry, routine diaper changes; linen only requires changing every 24 hours.					
FRICION-SHEAR Friction Occurs when Skin moves against support surfaces Shear Occurs when skin and adjacent bony surface slide across one another	1. Significant problem: Spasticity, contracture, itching, or agitation leads to almost constant thrashing and friction.	2. Problem: Requires moderate to maximum assistance in moving. Complete lifting without sliding against sheets is impossible. Frequently slides down in bed or chair, requiring frequent repositioning with maximum assistance.	3. Potential problem: Moves freely or requires minimum assistance. During a move, skin probably slides to some extent against sheets, chair, restraints, or other devices. Maintains relative good position in chair or bed most of the time but occasionally slides down.	4. No apparent problem: Able to completely lift patient during position change, moves in bed and in chair independently and has sufficient muscle strength to life up completely during move. Maintains good position in bed or chair at all times.*					
Nutritional Usual food intake pattern	1. Very Poor: NPO/or maintained on clear liquids, or IVs for more than 5 days OR albumin < 2.5 mg/dl OR never eats a complete meal. Rarely eats more than half of any food offered. Protein intake includes only 2 servings or meat or dairy products per day. Takes fluids poorly. Does not take a liquid dietary supplement.	2. Inadequate: Is on liquid diet or tube feedings/TPN, which provides inadequate calories and minerals for age OR albumin < 3 mg/dl OR rarely eats a complete meal and generally eats only about half of any food offered. Protein intake includes only 3 servings of meat or dairy products per day. Occasionally will take a dietary supplement.	3. Adequate: Is on tube feedings or TPN, which provide adequate calories and minerals for age OR eats over half of most meals. Eats a total of 4 servings of protein (meat, dairy products) each day. Occasionally will refuse a meal, but will usually take a supplement if offered.	4. Excellent: Is on a normal diet providing adequate calories for age. For example, eats most of every meal. Never refuses a meal. Usually eats a total of 4 or more servings of meat and dairy products. Occasionally eats between meals. Does not require supplementation.					
Tissue Perfusion & Oxygenation	1. Extremely compromised: Hypotensive (MAP < 50 mm Hg; < 40 in a newborn) or the patient does not physiologically tolerate position changes.	2. Compromised: Normotensive oxygen saturation may be < 95%; hemoglobin may be < 10 mg/dl; capillary refill may be > 2 seconds; serum pH is < 7.40.	3. Adequate: Normotensive oxygen saturation may be > 95%; hemoglobin may be < 10 mg/dl; capillary refill may be 2 seconds; serum pH is normal.	4. Excellent: Normotensive, oxygen saturation > 95%; normal hgb; capillary refill < 2 seconds.					
					TOTAL SCORE				
					Evaluator's Name				

Severe Risk : less than 9 | High Risk : 10-12 | Moderate Risk : 13-14 | Mild Risk : 15-18 | Not at Risk: 19-23

Risk Score	Category	Action	Support Surfaces (Please Note: Only required for children who are deemed at risk due to altered mobility, consider occupation therapy referral for advice)
15-18	At Risk	<ul style="list-style-type: none"> • Regular Turning Schedule • Enable as much activity as possible • Protect the heels • Use pressure redistribution surfaces • Manage moisture, friction and shear • Advance to a higher level of risk if other major risk factors are present 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
13-14	Moderate Risk	<ul style="list-style-type: none"> • Use the Same Protocol as for “At Risk” Patients • Position patient at 30 degree lateral incline using foam wedges 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
10-12	High Risk	<ul style="list-style-type: none"> • Follow the same protocol as for “Moderate Risk” Patients • In addition to regular turning schedule • Make small shifts in their position frequently 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
Less than 9	Severe Risk	<ul style="list-style-type: none"> • Use same protocol as for “High Risk” Patients • Add a pressure redistribution surface for patients with severe pain or with additional risk factors. 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay

Patient ID _____

BRADEN 'Q' SCALE



					Date :				
					Time :				
Mobility	1. Completely immobile: Does not make even slight changes in body or extremity position without assistance.	2. Very limited: Makes occasional slight changes in body or extremity position but unable to completely turn self independently.	3. Slightly limited: Makes frequent through slight changes in body or extremity position independently.	4. No limitations: Makes major and frequent changes in position without assistance.					
"Activity The degree of physical activity"	1. Bedfast : Confined to bed	2. Chairfast : Ability to walk severely limited or non-existent. Cannot bear own weight and/or must be assisted into chair or wheelchair."	3. Walks occasionally: Walks occasionally during day, but for very short distances, with or without assistance. Spends majority of each shift in bed or chair.	4. All patients too young to ambulate; OR walks frequently: Walks outside the room at least twice a day and inside room at least once every 2 hours during walking hours.					
Sensory Perception	1. Completely limited: Unresponsive (does not moan, flinch or grasp) to painful stimuli due to diminished level of consciousness or sedation, OR, limited ability to feel pain over most of the body surface.	2. Very limited: responds to only painful stimuli, cannot communicate discomfort except by moaning or restlessness; OR, has sensory impairment that limits the ability to feel pain or discomfort over half of body.	3. Slightly limited: Responds to verbal commands, but cannot always communicate discomfort or need to be turned; OR, has some sensory impairment that limits ability to feel pain, or discomfort in one or two extremities.	4. No impairment: Responds to verbal commands. Has no sensory deficit that would limit ability to feel or communicate pain or discomfort.					
Moisture Degree to which skin is exposed to moisture	1. Constantly moist: Skin is kept moist almost constantly by perspiration, urine, drainage, etc. Dampness is detected every time patient is moved or turned.	2. Very moist: Skin is often, but not always, moist. Linen must be changed at least every 8 hours.	3. Occasionally moist: Skin is occasionally moist, requiring linen change every 12 hours.	4. Rarely moist: Skin is usually dry, routine diaper changes; linen only requires changing every 24 hours.					
FRICTION-SHEAR Friction Occurs when Skin moves against support surfaces Shear Occurs when skin and adjacent bony surface slide across one another	1. Significant problem: Spasticity, contracture, itching, or agitation leads to almost constant thrashing and friction.	2. Problem: Requires moderate to maximum assistance in moving. Complete lifting without sliding against sheets is impossible. Frequently slides down in bed or chair, requiring frequent repositioning with maximum assistance.	3. Potential problem: Moves freely or requires minimum assistance. During a move, skin probably slides to some extent against sheets, chair, restraints, or other devices. Maintains relative good position in chair or bed most of the time but occasionally slides down.	4. No apparent problem: Able to completely lift patient during position change, moves in bed and in chair independently and has sufficient muscle strength to life up completely during move. Maintains good position in bed or chair at all times."					
Nutritional Usual food intake pattern	1. Very Poor: NPO/or maintained on clear liquids, or IVs for more than 5 days OR albumin < 2.5 mg/dl OR never eats a complete meal. Rarely eats more than half of any food offered. Protein intake includes only 2 servings of meat or dairy products per day. Takes fluids poorly. Does not take a liquid dietary supplement.	2. Inadequate: Is on liquid diet or tube feedings/TPN, which provides inadequate calories and minerals for age OR albumin < 3 mg/dl OR rarely eats a complete meal and generally eats only about half of any food offered. Protein intake includes only 3 servings of meat or dairy products per day. Occasionally will take a dietary supplement.	3. Adequate: Is on tube feedings or TPN, which provide adequate calories and minerals for age OR eats over half of most meals. Eats a total of 4 servings of protein (meat, dairy products) each day. Occasionally will refuse a meal, but will usually take a supplement if offered.	4. Excellent: Is on a normal diet providing adequate calories for age. For example, eats most of every meal. Never refuses a meal. Usually eats a total of 4 or more servings of meat and dairy products. Occasionally eats between meals. Does not require supplementation.					
Tissue Perfusion & Oxygenation	1. Extremely compromised: Hypotensive (MAP < 50 mm Hg; < 40 in a newborn) or the patient does not physiologically tolerate position changes.	2. Compromised: Normotensive oxygen saturation may be < 95%; hemoglobin may be < 10 mg/dl; capillary refill may be > 2 seconds; serum pH is < 7.40.	3. Adequate: Normotensive oxygen saturation may be < 95%; hemoglobin may be < 10 mg/dl; capillary refill may be 2 seconds; serum pH is normal.	4. Excellent: Normotensive, oxygen saturation > 95%; normal hgb; capillary refill < 2 seconds.					

Severe Risk : less than 9 | High Risk : 10-12 | Moderate Risk : 13-14 | Mild Risk : 15-18 | Not at Risk: 19-23

Docu. No. : RCH /FRM / CLINICAL / 119

TOTAL SCORE

Evaluator's Name

Risk Score	Category	Action	Support Surfaces (Please Note: Only required for children who are deemed at risk due to altered mobility, consider occupation therapy referral for advice)
15-18	At Risk	<ul style="list-style-type: none"> • Regular Turning Schedule • Enable as much activity as possible • Protect the heels • Use pressure redistribution surfaces • Manage moisture, friction and shear • Advance to a higher level of risk if other major risk factors are present 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
13-14	Moderate Risk	<ul style="list-style-type: none"> • Use the Same Protocol as for “At Risk” Patients • Position patient at 30 degree lateral incline using foam wedges 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
10-12	High Risk	<ul style="list-style-type: none"> • Follow the same protocol as for “Moderate Risk” Patients • In addition to regular turning schedule • Make small shifts in their position frequently 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
Less than 9	Severe Risk	<ul style="list-style-type: none"> • Use same protocol as for “High Risk” Patients • Add a pressure redistribution surface for patients with severe pain or with additional risk factors. 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay

Patient ID _____

BRADEN 'Q' SCALE



					Date :				
					Time :				
Mobility	1. Completely immobile: Does not make even slight changes in body or extremity position without assistance.	2. Very limited: Makes occasional slight changes in body or extremity position but unable to completely turn self independently.	3. Slightly limited: Makes frequent through slight changes in body or extremity position independently.	4. No limitations: Makes major and frequent changes in position without assistance.					
Activity The degree of physical activity	1. Bedfast : Confined to bed	2. Chairfast : Ability to walk severely limited or non-existent. Cannot bear own weight and/or must be assisted into chair or wheelchair.*	3. Walks occasionally: Walks occasionally during day, but for very short distances, with or without assistance. Spends majority of each shift in bed or chair.	4. All patients too young to ambulate; OR walks frequently: Walks outside the room at least twice a day and inside room at least once every 2 hours during walking hours.					
Sensory Perception	1. Completely limited: Unresponsive (does not moan, flinch or grasp) to painful stimuli due to diminished level of consciousness or sedation. OR, limited ability to feel pain over most of the body surface.	2. Very limited: responds to only painful stimuli, cannot communicate discomfort except by moaning or restlessness; OR, has sensory impairment that limits the ability to feel pain or discomfort over half of body.	3. Slightly limited: Responds to verbal commands, but cannot always communicate discomfort or need to be turned; OR, has some sensory impairment that limits ability to feel pain, or discomfort in one or two extremities.	4. No impairment: Responds to verbal commands. Has no sensory deficit that would limit ability to feel or communicate pain or discomfort.					
Moisture Degree to which skin is exposed to moisture	1. Constantly moist: Skin is kept moist almost constantly by perspiration, urine, drainage, etc. Dampness is detected every time patient is moved or turned.	2. Very moist: Skin is often, but not always, moist. Linen must be changed at least every 8 hours.	3. Occasionally moist: Skin is occasionally moist, requiring linen change every 12 hours.	4. Rarely moist: Skin is usually dry, routine diaper changes; linen only requires changing every 24 hours.					
FRICION-SHEAR Friction Occurs when Skin moves against support surfaces Shear Occurs when skin and adjacent bony surface slide across one another	1. Significant problem: Spasticity, contracture, itching, or agitation leads to almost constant thrashing and friction.	2. Problem: Requires moderate to maximum assistance in moving. Complete lifting without sliding against sheets is impossible. Frequently slides down in bed or chair, requiring frequent repositioning with maximum assistance.	3. Potential problem: Moves freely or requires minimum assistance. During a move, skin probably slides to some extent against sheets, chair, restraints, or other devices. Maintains relative good position in chair or bed most of the time but occasionally slides down.	4. No apparent problem: Able to completely lift patient during position change, moves in bed and in chair independently and has sufficient muscle strength to life up completely during move. Maintains good position in bed or chair at all times.*					
Nutritional Usual food intake pattern	1. Very Poor: NPO/or maintained on clear liquids, or IVs for more than 5 days OR albumin < 2.5 mg/dl OR never eats a complete meal. Rarely eats more than half of any food offered. Protein intake includes only 2 servings of meat or dairy products per day. Takes fluids poorly. Does not take a liquid dietary supplement.	2. Inadequate: Is on liquid diet or tube feedings/TPN, which provides inadequate calories and minerals for age OR albumin < 3 mg/dl OR rarely eats a complete meal and generally eats only about half of any food offered. Protein intake includes only 3 servings of meat or dairy products per day. Occasionally will take a dietary supplement.	3. Adequate: Is on tube feedings or TPN, which provides adequate calories and minerals for age OR eats over half of most meals. Eats a total of 4 servings of protein (meat, dairy products) each day. Occasionally will refuse a meal, but will usually take a supplement if offered.	4. Excellent: is on a normal diet providing adequate calories for age. For example, eats most of every meal. Never refuses a meal. Usually eats a total of 4 or more servings of meat and dairy products. Occasionally eats between meals. Does not require supplementation.					
Tissue Perfusion & Oxygenation	1. Extremely compromised: Hypotensive (MAP < 50 mm Hg; < 40 in a newborn) or the patient does not physiologically tolerate position changes.	2. Compromised: Normotensive oxygen saturation may be < 95%; hemoglobin may be < 10 mg/dl; capillary refill may be > 2 seconds; serum pH is < 7.40.	3. Adequate: Normotensive oxygen saturation may be < 95%; hemoglobin may be < 10 mg/dl; capillary refill may be 2 seconds; serum pH is normal.	4. Excellent: Normotensive, oxygen saturation > 95%; normal hgb; capillary refill < 2 seconds.					

Severe Risk : less than 9 | High Risk : 10-12 | Moderate Risk : 13-14 | Mild Risk : 15-18 | Not at Risk: 19-23

Docu. No. : RCH / FRM / CLINICAL / 119

TOTAL SCORE

Evaluator's Name

Risk Score	Category	Action	Support Surfaces (Please Note: Only required for children who are deemed at risk due to altered mobility, consider occupation therapy referral for advice)
15-18	At Risk	<ul style="list-style-type: none"> • Regular Turning Schedule • Enable as much activity as possible • Protect the heels • Use pressure redistribution surfaces • Manage moisture, friction and shear • Advance to a higher level of risk if other major risk factors are present 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
13-14	Moderate Risk	<ul style="list-style-type: none"> • Use the Same Protocol as for "At Risk" Patients • Position patient at 30 degree lateral incline using foam wedges 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
10-12	High Risk	<ul style="list-style-type: none"> • Follow the same protocol as for "Moderate Risk" Patients • In addition to regular turning schedule • Make small shifts in their position frequently 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay
Less than 9	Severe Risk	<ul style="list-style-type: none"> • Use same protocol as for "High Risk" Patients • Add a pressure redistribution surface for patients with severe pain or with additional risk factors. 	High density foam mattress Gel pads for high-risk areas Alternating pressure mattress overlay

LBM-00112798 IP26-00006647
 Master ARJUN KARTHIKEYA
 13-02-2025 1 Y 4 M 13 D (M)
 Dr. BINDHURA MUNUKUNTALA



Morse Fall Risk Assessment Form

Choose Highest Applicable Score from each Category		Date / Time			Fall Risk Grading		
		Score			Risk Level	Morse Fall Score (MFS)	Action
History of Falling (immediately or w/in 3 months)	Yes	25			Low Risk	0 - 24	Standard Fall Precaution
	No	0					
Secondary Diagnosis (more than one diagnosis)	Yes	15			Moderate Risk	25 - 50	Implement Moderate Fall Prevention Intervention
	No	0					
Ambulatory Aid	Furniture	30			High Risk	≥ 51	Implement High Risk Fall Prevention Intervention
	Crutches, Cane(S), Walker	15					
	None /Bed Rest /Nurse Assist	0					
IV / Heparin Lock or Saline	Yes	20			Moderate Risk	25 - 50	Implement Moderate Fall Prevention Intervention
	No	0					
GAIT / Transferring	Impaired	20			High Risk	≥ 51	Implement High Risk Fall Prevention Intervention
	Weak (uses touch for balance)	10					
	Normal /On Bed Rest /Immobile	0					
Mental Status	Forgets limitations	15			High Risk	≥ 51	Implement High Risk Fall Prevention Intervention
	Oriented to own ability	0					
Total Morse Fall Scale Score:							
		Signature					

Tick (✓) whichever precaution taken. ✓

Risk Level and Interventions

Low Risk (0 - 24) (Standard Falls Precautions)

- Ensure patients use their prescribed eye glasses if any, in the hospital
- Use chairs with arm rests
- Use safety straps on stretchers and wheelchairs while transporting patients

Moderate Risk (25-50) Apply all low risk intervention and

- Assist and/or supervise ambulation. Reinforce to always call for assistance
- Hourly safety check
- Assess patient after visitors, leave to ensure safety measures in place

High Risk (≥ 51) Apply all low and moderate risk interventions, and.

- Initiate constant observation by healthcare provider as appropriate to patient's needs

Morse Fall Risk Assessment Form

Choose Highest Applicable Score from each Category		Date / Time			Fall Risk Grading		
		Score			Risk Level	Morse Fall Score (MFS)	Action
History of Falling (immediately or w/in 3 months)	Yes	25			Low Risk	0 - 24	Standard Fall Precaution
	No	0					
Secondary Diagnosis (more than one diagnosis)	Yes	15			Moderate Risk	25 - 50	Implement Moderate Fall Prevention Intervention
	No	0					
Ambulatory Aid	Furniture	30			High Risk	≥ 51	Implement High Risk Fall Prevention Intervention
	Crutches, Cane(S), Walker	15					
	None /Bed Rest /Nurse Assist	0					
IV / Heparin Lock or Saline	Yes	20			Moderate Risk	25 - 50	Implement Moderate Fall Prevention Intervention
	No	0					
GAIT / Transferring	Impaired	20			High Risk	≥ 51	Implement High Risk Fall Prevention Intervention
	Weak (uses touch for balance)	10					
	Normal /On Bed Rest /Immobile	0					
Mental Status	Forgets limitations	15			High Risk	≥ 51	Implement High Risk Fall Prevention Intervention
	Oriented to own ability	0					
Total Morse Fall Scale Score:							
		Signature					

Tick (✓) whichever precaution taken.

Risk Level and Interventions

Low Risk (0 – 24) (Standard Falls Precautions)

- Ensure patients use their prescribed eye glasses if any, in the hospital
- Use chairs with arm rests
- Use safety straps on stretchers and wheelchairs while transporting patients

Moderate Risk (25-50) Apply all low risk intervention and

- Assist and/or supervise ambulation. Reinforce to always call for assistance
- Hourly safety check
- Assess patient after visitors, leave to ensure safety measures in place

High Risk (≥ 51) Apply all low and moderate risk interventions, and.

- Initiate constant observation by healthcare provider as appropriate to patient's needs

Morse Fall Risk Assessment Form

Choose Highest Applicable Score from each Category		Date / Time			Fall Risk Grading		
		Score			Risk Level	Morse Fall Score (MFS)	Action
History of Falling (Immediately or w/in 3 months)	Yes	25			Low Risk	0 - 24	Standard Fall Precaution
	No	0					
Secondary Diagnosis (more than one diagnosis)	Yes	15			Moderate Risk	25 - 50	Implement Moderate Fall Prevention Intervention
	No	0					
Ambulatory Aid	Furniture	30			High Risk	≥ 51	Implement High Risk Fall Prevention Intervention
	Crutches, Cane(S), Walker	15					
	None /Bed Rest /Nurse Assist	0					
IV / Heparin Lock or Saline	Yes	20			Moderate Risk	25 - 50	Implement Moderate Fall Prevention Intervention
	No	0					
GAIT / Transferring	Impaired	20			High Risk	≥ 51	Implement High Risk Fall Prevention Intervention
	Weak (uses touch for balance)	10					
	Normal /On Bed Rest /Immobile	0					
Mental Status	Forgets limitations	15			High Risk	≥ 51	Implement High Risk Fall Prevention Intervention
	Oriented to own ability	0					
Total Morse Fall Scale Score:							
		Signature					

Tick (✓) whichever precaution taken.

Risk Level and Interventions

Low Risk (0 – 24) (Standard Falls Precautions)

- Ensure patients use their prescribed eye glasses if any, in the hospital
- Use chairs with arm rests
- Use safety straps on stretchers and wheelchairs while transporting patients

Moderate Risk (25-50) Apply all low risk intervention and

- Assist and/or supervise ambulation. Reinforce to always call for assistance
- Hourly safety check
- Assess patient after visitors, leave to ensure safety measures in place

High Risk (≥ 51) Apply all low and moderate risk interventions, and.

- Initiate constant observation by healthcare provider as appropriate to patient's needs



CHECKLIST FOR THROMBOPHLEBITIS

S. No.	SITE OBSERVATION	STAGE / ACTION	SCORE	DAY-1/6			DAY-2			DAY-3			Remarks
				M	E	N	M	E	N	M	E	N	
1	IV site appears healthy	No signs of phlebitis / Observe cannula	0			0							
2	One of the following signs is evident : * Slight pain near the IV Site / * Slight redness near IV Site	Possibly first signs of phlebitis / Observe cannula	1			0							
3	Two of the following Signs are evident: Pain at IV site Redness	Early stage of phlebitis / Resite Cannula	2			0							
4	All of the following Signs are evident : Pain along Path of cannula Redness around Site Swelling	Medium stage of phlebitis / Resite Cannula Consider Treatment	3			0							
5	All of the following Signs are evident and Extensive : Pain along Path of cannula Redness around Site Swelling palpable Venous cord	Advanced stage of phlebitis or the start of thrombophlebitis / Re site Cannula Consider Treatment	4			0							
6	All of the following Signs are evident and Extensive : Pain along Path of cannula Redness around Site Swelling palpable Venous cord pyrexia	Advanced stage of thrombophlebitis / Initiate treatment Re site Cannula	5			0							
Signature of the Nurse						CS							

NOTE : Phlebitis greater than grade 2 should be reported to physicians and other appropriate health care personal ongoing observation of the site should continue for 48 hours post removal to detect post infusion phlebitis.

Signature of Shift In Charge :

Signature : Name :

Signature of Ward In Charge :

Signature : Name :

CHECKLIST FOR THROMBOPHLEBITIS

S. No.	SITE OBSERVATION	STAGE / ACTION	SCORE	DAY-1			DAY-2			DAY-3			Remarks
				M	E	N	M	E	N	M	E	N	
1	IV site appears healthy	No signs of phlebitis / Observe cannula	0										
2	One of the following signs is evident : * Slight pain near the IV Site / * Slight redness near IV Site	Possibly first signs of phlebitis / Observe cannula	1										
3	Two of the following Signs are evident: Pain at IV site Redness	Early stage of phlebitis / Resite Cannula	2										
4	All of the following Signs are evident : Pain along Path of cannula Redness around Site Swelling	Medium stage of phlebitis / Resite Cannula Consider Treatment	3										
5	All of the following Signs are evident and Extensive : Pain along Path of cannula Redness around Site Swelling palpable Venous cord	Advanced stage of phlebitis or the start of thrombophlebitis / Re site Cannula Consider Treatment	4										
6	All of the following Signs are evident and Extensive : Pain along Path of cannula Redness around Site Swelling palpable Venous cord pyrexia	Advanced stage of thrombophlebitis / Initiate treatment Re site Cannula	5										
Signature of the Nurse													

NOTE : Phlebitis greater than grade 2 should be reported to physicians and other appropriate health care personal ongoing observation of the site should continue for 48 hours post removal to detect post infusion phlebitis.

Signature of Shift In Charge :

Signature of Ward In Charge :

Signature : Name :

Signature : Name :

NURSING SHIFT HAND OVER FORM

SITUATION	Diagnosis:	Any Infection: <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not Known If Yes Specify:						
	Surgery / Procedure:	Post OP Day:						
BACKGROUND	Date	25/6/25						
	Shift	ICU						
	Medical Condition (Any special condition to be noted):	-						
	Diet:	-						
ASSESSMENT	Allergy:	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	Ventilation (RA, NP, NIV, VENTI):	-						
	Tubes/Drains/Catheter:	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	Vital Signs:	Temp:	96.5 F					
		Res:	20b/m					
		SpO ₂ :	99%					
		Pulse:	102b/m					
		BP:	-					
		LOC:	-					
		Fall Risk Score:	-					
Pain Score:	-							
Skin Integrity	-							
Recommendations	Safety Needs:	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	Physiotherapy:	-						
	Others Specify:	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	Special Diet:	-						
	Critical Lab Test / Values:	-						
	Other Special Orders / Medications:	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	PU Prophylaxis:	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
DVT Prophylaxis:	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No		
ADL (Dependent / Non Dependent):	-							
Post Operative Procedure Special Orders:		NA						
Handed Over By Name :		Alecobsh						
Signature / ID :		25/6/25						
Date:		SPM						
Time:								
Taken Over By Name :								
Signature / ID :								
Date:								
Time:								

Patient Sticker



NURSING SHIFT HAND OVER FORM

SITUATION	Diagnosis:	Any Infection: <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not Known If Yes Specify:						
	Surgery / Procedure:	Post OP Day:						
BACKGROUND	Date	Shift	/	/	/	/	/	
	Medical Condition (Any special condition to be noted):							
	Diet:							
ASSESSMENT	Allergy:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	Ventilation (RA, NP, NIV, VENTI):							
	Tubes/Drains/Catheter:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	Vital Signs:	Temp:						
		Res:						
		SpO ₂ :						
		Pulse:						
		BP:						
		LOC:						
		Fall Risk Score:						
Pain Score:								
Skin Integrity								
Recommendations	Safety Needs:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	Physiotherapy:							
	Others Specify:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	Special Diet:							
	Critical Lab Test / Values:							
	Other Special Orders / Medications:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
	ADL (Dependent / Non Dependent):							
PU Prophylaxis:		<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
DVT Prophylaxis:		<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	
Post Operative Procedure Special Orders:								
Handed Over By Name :								
Signature / ID :								
Date:								
Time:								
Taken Over By Name :								
Signature / ID :								
Date:								
Time:								

Patient Sticker



NURSING SHIFT HAND OVER FORM

SITUATION	Diagnosis:	Any Infection: <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not Known If Yes Specify:					
	Surgery / Procedure:	Post OP Day:					
BACKGROUND	Date	/	/	/	/	/	/
	Shift						
	Medical Condition (Any special condition to be noted):						
	Diet:						
ASSESSMENT	Allergy:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	Ventilation (RA, NP, NIV, VENTI):						
	Tubes/Drains/Catheter:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	Vital Signs:	Temp:					
		Res:					
		SpO ₂ :					
		Pulse:					
		BP:					
		LOC:					
		Fall Risk Score:					
	Pain Score:						
	Skin Integrity						
Recommendations	Safety Needs:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	Physiotherapy:						
	Others Specify:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	Special Diet:						
	Critical Lab Test / Values:						
	Other Special Orders / Medications:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	PU Prophylaxis:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	DVT Prophylaxis:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
ADL (Dependent / Non Dependent):							
Post Operative Procedure Special Orders:							
Handed Over By Name :							
Signature / ID :							
Date:							
Time:							
Taken Over By Name :							
Signature / ID :							
Date:							
Time:							

Patient Sticker



NURSING SHIFT HAND OVER FORM

SITUATION	Diagnosis:	Any Infection: <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Not Known If Yes Specify:					
	Surgery / Procedure:	Post OP Day:					
BACKGROUND	Date	Shift	/	/	/	/	/
	Medical Condition (Any special condition to be noted):						
	Diet:						
ASSESSMENT	Allergy:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	Ventilation (RA, NP, NIV, VENTI):						
	Tubes/Drains/Catheter:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	Vital Signs:	Temp:					
		Res:					
		SpO ₂ :					
		Pulse:					
		BP:					
		LOC:					
		Fall Risk Score:					
	Pain Score:						
	Skin Integrity						
Recommendations	Safety Needs:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	Physiotherapy:						
	Others Specify:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	Special Diet:						
	Critical Lab Test / Values:						
	Other Special Orders / Medications:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	ADL (Dependent / Non-Dependent):						
	PU Prophylaxis:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
	DVT Prophylaxis:	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Yes <input type="checkbox"/> No
Post Operative Procedure Special Orders:							
Handed Over By Name :							
Signature / ID :							
Date:							
Time:							
Taken Over By Name :							
Signature / ID :							
Date:							
Time:							